



METHODS FOR

“Trends in drug-related hospitalisations in Australia, 1999-2021”

**Agata Chrzanowska¹, Nicola Man¹, Jane Akhurst¹,
Rachel Sutherland¹, Louisa Degenhardt¹
and Amy Peacock^{1,2}**

¹ National Drug and Alcohol Research Centre, University of New South Wales

² School of Psychology, University of Tasmania

ISBN 978-0-7334-4058-8 ©NDARC, UNSW SYDNEY 2022

This report was prepared by researchers from the National Drug and Alcohol Research Centre for the Drug Trends program. The Drug Trends program is coordinated by the National Drug and Alcohol Research Centre, UNSW Sydney and undertaken in partnership with the Burnet Institute, National Drug Research Institute, University of Queensland, and University of Tasmania.

This work is copyright. You may download, display, print and reproduce this material in unaltered form only (retaining this notice) for your personal, non-commercial use or use within your organisation. All other rights are reserved. Requests and enquiries concerning reproduction and rights should be addressed to NDARC, UNSW Sydney, NSW 2052, Australia.

Recommended citation: Chrzanowska, A., Man, N., Akhurst, J., Sutherland, R., Degenhardt, L. & Peacock, A. (2022). Trends in drug-related hospitalisations in Australia, 1999-2021. Sydney: National Drug and Alcohol Research Centre, UNSW Sydney. [DOI: 10.26190/wrsv-3b78](https://doi.org/10.26190/wrsv-3b78)

Please note that as with all statistical reports there is the potential for minor revisions to data in this report over its life. Please refer to the online version at [Drug Trends](#).

Please contact the Drug Trends team with any queries regarding this publication:
drugtrends@unsw.edu.au.

Table of Contents

GENERAL NOTE	5
SCOPE OF THE DATA	5
PRESENTATION OF RESULTS	6
AGE AND SEX	7
JURISDICTION AND REMOTENESS AREA.....	7
LIMITATION IN JURISDICTIONAL DATA.....	8
TASMANIAN RECORDS	8
VICTORIAN RECORDS.....	8
TOTAL NUMBER OF HOSPITALISATIONS IN AUSTRALIAN CAPITAL TERRITORY, NORTHERN TERRITORY AND TASMANIA	8
TERMINOLOGY.....	8
CODING OF HOSPITALISATIONS.....	10
ALL DRUG-RELATED HOSPITALISATIONS.....	10
PRINCIPAL DIAGNOSIS AND EXTERNAL CAUSE	11
OPIOID-RELATED HOSPITALISATIONS	12
NON-OPIOID ANALGESIC-RELATED HOSPITALISATIONS	13
ANTI-EPILEPTIC, SEDATIVE-HYPNOTIC AND ANTIPARKINSONISM DRUG-RELATED HOSPITALISATIONS	14
CANNABINOID-RELATED HOSPITALISATIONS	15
HALLUCINOGEN-RELATED HOSPITAL SEPARATIONS.....	15
COCAINE-RELATED HOSPITALISATIONS	16
AMPHETAMINE-TYPE STIMULANT-RELATED HOSPITALISATIONS	16
ANTIDEPRESSANT, ANTIPSYCHOTIC AND NEUROLEPTIC-RELATED HOSPITALISATIONS	17
VOLATILE SOLVENT-RELATED HOSPITALISATIONS	18
MULTIPLE DRUG USE-RELATED HOSPITALISATIONS	19

Acknowledgements

Funding

The Drug Trends program is funded by the Australian Government Department of Health under the Drug and Alcohol Program.

Data source

We would like to acknowledge the Australian Institute of Health and Welfare for data from the National Hospital Morbidity Database.

Contributors

We would like to acknowledge the contribution of those who have been involved in past reporting on drug-related hospitalisations by Drug Trends, specifically: A/Prof Timothy Dobbins, Dr Amanda Roxburgh, and A/Prof Lucinda Burns.

We thank Dr Louise Tierney and her team from the Tobacco, Alcohol and Other Drugs Unit at the Australian Institute of Health and Welfare for reviewing the report.

We acknowledge the traditional custodians of the land on which the work for this report was undertaken. We pay respect to Elders past, present, and emerging.

Related Links

- Hospitalisations data visualisations: https://drugtrends.shinyapps.io/hospital_separations
- Hospitalisations methods document: <https://ndarc.med.unsw.edu.au/resource-analytics/trends-drug-related-hospitalisations-australia-1999-2021>
- For other Drug Trends publications on drug-related hospitalisations and drug-induced deaths in Australia, go to: <https://ndarc.med.unsw.edu.au/project/national-illicit-drug-indicators-project-nidip>
- For more information on NDARC research, go to: <http://ndarc.med.unsw.edu.au/>
- For more information about the AIHW and NHMD, go to: <https://www.aihw.gov.au/>
- For more information on ICD coding go to: <http://www.who.int/classifications/icd/en/>
<https://www.ihacpa.gov.au/resources/icd-10-amachiacs-eleventh-edition>
- For more research from the Drug Trends program go to: <https://ndarc.med.unsw.edu.au/program/drug-trends>

General note

Data on hospitalisations presented in the [online interactive visualisation](#) and reported in the [report](#) were derived from the [National Hospital Morbidity Database](#) (NHMD) held by the Australian Institute of Health and Welfare (AIHW). Ethical approval to obtain the data, analyse them and present in this report was provided by the UNSW Human Research Ethics Committee.

Scope of the data

The National Hospital Morbidity Database (NHMD) is compiled from data supplied by the state and territory health authorities. It is a collection of electronic confidentialised summary records for hospital [separations](#) (that is, completed [episodes of care](#)) in public and private hospitals in Australia. Separations for which the care type was reported as 'newborn without qualified days', and records for '[posthumous organ procurement](#)' and '[hospital boarders](#)' have been excluded. It should be noted that state of hospitalisation equals the state of usual residence, and that cross-border separations were not provided. Hospitalisations in Western Australia with a contracted patient status of 'Inter-hospital contracted patient to private sector hospital', were also not provided to adjust for separations recorded on both sides of contractual care arrangements.

Although there are national standards for data on hospital services, there are some variations in how hospital services are defined and counted, between public and private hospitals, among the states and territories and over time. The national data on hospital care does not include care provided by non-hospital providers, such as community health centres. For more information, see the [AIHW report](#) 'Variation in hospital admission policies and practices: Australian hospital statistics'.

At the time of separation, a principal (i.e., main) diagnosis, and up to 99 additional diagnoses may be recorded. The data presented in this report include only hospitalisations where drugs were determined to be the principal reason for the hospital admission (i.e., identified in the principal diagnosis field).

We have not included hospitalisations where the principal diagnosis is related to tobacco or alcohol use, other unspecified drug use and fetal and perinatal conditions. It is important to note many drug-related hospitalisations involve more than one drug (including alcohol) but may have one substance coded as the 'principal diagnosis'. Further, sometimes it is not possible to determine one substance as the primary drug leading to the hospitalisation; these cases are coded and presented as 'multiple drug use' and thus will not be represented in the count of hospitalisations for a single substance.

Hospitalisations are coded according to the World Health Organization's (WHO) International Statistical Classification of Diseases (ICD) and Related Problems, [Australian Modification](#). The ICD 10th revision (ICD-10-AM) (National Centre for Classification in Health, 1998) was used to code data dating from 1999 to the present in South Australia (SA), Western Australia (WA), and Queensland (QLD). The remaining jurisdictions commenced using ICD-10-AM codes in 1998. Caution should be exercised in comparing diagnosis, procedure and external cause data over time, as the classifications and coding standards for these data can change. Drug involvement in presentation to hospital may be underreported or under-ascertained, and the coding system used does not necessarily allow identification of the specific drug involved in presentations (often instead the broader substance category e.g., 'synthetic opioids' rather than 'fentanyl'). Please refer to Chronicle of [ICD-10-AM/ACHI/ACS](#) First-Eleventh Edition for information on changes in classifications and coding standards over time and ICD-10-AM editions.

Due to the different ways in which diagnoses are classified in the 9th and 10th revision of the ICD coding structure, analyses presented in this report include data from 1999-2000 to 2020-21 only. For trends in drug-related hospitalisations from 1993-94 to 2014-15 on a restricted set of diagnosis and/or drugs, please refer to our [previous reports](#) or contact the authors (drugtrends@unsw.edu.au).

Presentation of results

Results of the analysis of the hospitalisations data are presented as:

- **number of hospitalisations**,
- **percentage**,
- **crude rate of hospitalisations per 100,000 people** with corresponding 95% confidence intervals (95% CI), calculated using the Australian Bureau of Statistics [estimated resident population](#) (ERP) figures (including population by sex, age group, remoteness and jurisdiction) as of 30 June each year released on 28th June 2022, and
- **age-standardised rate per 100,000 people** with corresponding 95% CI, calculated using the Australian Bureau of Statistics [ERP](#) figures as of 30 June each year released on 28th June 2022 and the [Australian Standard Population](#) at 30 June 2001.

Where we discuss changes in rate of drug-related hospitalisations during the COVID-19 pandemic, we report quarterly rates calculated as population crude rates per 100,000 estimated resident population (ERP) using [quarterly ERPs](#). It should be noted that there has been disruption in the usual population trends caused by the closure of the international borders from 20 March 2020. In the financial year July 2020 to June 2021, the overall [population growth](#) was much smaller than the years prior, hence caution is required when interpreting statistics calculated from the ERPs and comparing them over time ([National, state and territory population, June 2021, Australian Bureau of Statistics \(abs.gov.au\)](#)).

Estimates from data with counts less than or equal to 5 have been suppressed to protect confidentiality. Age-standardised rates enable the comparison of rates over time and between populations of different age structures. Age-standardised rates were calculated using the [direct](#) standardisation method as recommended by the Australian Bureau of Statistics (ABS) and AIHW. In accordance with recommendations to ensure stability of age-standardised rates from sparse data, age-standardised rates were not calculated if the number of total hospitalisations was less than or equal to 10. In this case, the reader should refer to other measures such as number of deaths or crude rate of deaths. Age-standardised rates are only calculated for people of all ages (i.e., the rate is not calculated for specific age groups). In the [online interactive visualisation](#), data are not displayed if the age-standardised rate is selected for age group analysis.

Crude rate per 100,000 population and age-standardised rate per 100,000 population are computed of the population of interest (e.g., rate among females is computed of the female population).

Rate ratio (crude or age-standardised; *RR*) was calculated to compare the rate for the year 2020-21 ($rate_{2020-21}$) with the corresponding rate for the year 2019-20 ($rate_{2019-20}$) as follows:

$$RR = \frac{rate_{2020-21}}{rate_{2019-20}}$$

The 95% confidence interval of the crude rate ratio is calculated using the `ir` command in Stata version 16.0 which uses the exact binomial method described by Rothman (1986).

The standard error for the log of the age-standardised rate ratio, $SE(\ln(RR))$, given the variance of the age-standardised rate in 2020-21 ($var(rate_{2020-21})$) and 2019-20 ($var(rate_{2019-20})$), was firstly calculated as:

$$SE(\ln(RR)) = \sqrt{\frac{var(rate_{2020-21})}{rate_{2020-21}^2} + \frac{var(rate_{2019-20})}{rate_{2019-20}^2}}$$

The 95% confidence interval of the age-standardised rate ratio $95\%CI(RR)$ was calculated as:

$$95\%CI(RR) = \exp(\ln(RR) \pm 1.96 \times SE(\ln(RR))) \quad \text{Rothman (1986)}$$

Rate ratios range between 0 and infinity. In our report, a rate ratio of <1 indicates that there was a decrease in the rate of hospitalisations from 2019-20 to 2020-21 and a rate ratio of >1 indicates that there was an increase in the rate of hospitalisations from 2019-20 to 2020-21. The threshold for statistical significance is $p < 0.05$.

Age and sex

Age of the person at the time of admission to hospital is provided by the AIHW in the NHMD as a 5-year and 10-year age group. In the report and in the online data visualisation we report on findings for Australians of all ages and by 10-year age group (10-19 to 60-69 and 70 and over), where data allows for such disaggregation. We do not report on finding for age under 10 years due to sensitivity of these data, however this age group is included in analysis of total hospitalisations. [Sex](#) is reported as male and female, as provided by the AIHW in the NHMD. Hospitalisations where sex was reported as intersex or indeterminate or was not stated/inadequately described are grouped in the NHMD as one category. These records are not included in sex analysis but are included in analyses of total hospitalisations.

Jurisdiction and remoteness area

Data on state and territory is defined by the place of usual residence of the admitted patient. Records where the state of hospitalisation was the same as the state of usual residence were provided (i.e., cross-border separations were excluded).

Data on the remoteness area of usual residence was classified, using the Australian Statistical Geography Standard ([ASGS](#)), into five categories: Major Cities, Inner Regional, Outer Regional, Remote, and Very Remote Australia. These data have been collected by AIHW in the NHMD dataset since 2012-13. Due to small numbers, we further combined Remote and Very Remote Australia into one category to protect confidentiality. Data on remoteness was released by all jurisdictions, except Queensland, until 2018-19. The omission of Queensland data means that national trend analysis could not be presented for the complete time period and only summary statistics and change between 2018-19 and 2020-21 are available for Queensland and Australia. Trend analysis for period 2012-13 to 2020-21 are available for other jurisdictions for which data were released.

Limitation in jurisdictional data

Tasmanian records

For Tasmanian records, provision of data between 2008-09 and 2015-16 was limited to selected drug and alcohol related principal and/or additional diagnoses and external causes. Specifically, of the ICD-10-AM codes specified above, only records with following codes were provided:

- Principal diagnosis codes at full character level (where applicable) for F10, F11, F12, F13, F14, F15, F16, F19, T40, T423, T424, T427, T436, T439.

Particularly, many of the T-codes related to poisoning are missing. Estimates of drug-related hospitalisations for this period are likely to be underestimated as a consequence.

Victorian records

From 1st July 2011 to 30th June 2013 (i.e., between 2011-12 and 2012-13), there was a large decrease in public hospitalisations reported for the Victorian Admitted Episodes Dataset (VAED) because episodes where the patient's entire care is provided in the emergency department were not considered for admission, irrespective of whether a criterion for admission is met. From 2013-14 onwards, "ED-only admissions" were largely replaced with admissions to Short Stay Observation Units.

Total number of hospitalisations in Australian Capital Territory, Northern Territory and Tasmania

Total number of hospitalisations in public and private hospitals in 2020-21 was not available for Australian Capital Territory, Northern Territory and Tasmania. For those jurisdictions only data on the number of hospitalisations from public hospitals was available in the [Admitted patients](#) report by the Australian Institute of Health and Welfare, hence the percentage of drug-related hospitalisation of all hospitalisations in these jurisdictions could not be calculated.

Terminology

Terminology used in this report aligns with terminology used by AIHW in the report [Drug related hospitalisations](#) to describe findings from the NHMD.

An [admitted patient](#) is a patient who undergoes a hospital's formal admission process to receive treatment and/or care.

A **hospitalisation** in our reporting (also called hospital [separation](#)) refers to a completed [episode of admitted patient's care](#) in a hospital ending with discharge, death, transfer or a portion of a hospital stay beginning or ending in a change to another type of care. There can be more than one hospitalisation for each patient and hospitalisation can be either same-day (hospital admission and separation happen on the same day) or overnight (hospital admission and separation happen on a different date).

Each NHMD separation record includes one principal diagnosis and up to 99 additional diagnoses.

The [principal diagnosis](#) is defined as the diagnosis determined after study and established at the completion of the episode of care to be chiefly responsible for occasioning the patient's episode of admitted patient care.

Additional diagnoses are conditions or complaints that either coexists with the principal diagnosis or develop during the episode of care and affect patient's management.

An **external cause** is defined as the event, circumstance or condition associated with the occurrence of injury, poisoning or violence. Whenever a patient has a principal or additional diagnosis of an injury or poisoning, an external cause should be recorded.

A **drug-related hospitalisation** refers to hospital care with principal diagnosis of selected substance-use disorder or harm due to selected substances. Hospitalisations where the diagnosis of drug-related harm or disorder is additional to the principal diagnosis, such as problems related to certain chronic conditions, have been excluded and, as aforementioned, hospitalisations where alcohol or tobacco comprise the principal diagnosis are not included.

Coding of hospitalisations

All drug-related hospitalisations

The following ICD-10-AM codes were used to examine trends in drug-related hospitalisations from 1999-00 to 2020-21.

Drug type	Examples of drugs commonly assigned to ICD-10-AM category	ICD-10-AM
Opioids		F11.0–F11.9, T40.0–T40.4, T40.6
Opium		T40.0
Heroin		T40.1
Natural and semi-synthetic opioids	Oxycodone, Morphine, Codeine	T40.2
Methadone		T40.3
Synthetic opioids	Fentanyl, Tramadol, Pethidine	T40.4
Other and unspecified opioids		T40.6
Non-opioid analgesics	Paracetamol, Ibuprofen, Aspirin	F55.2, T39.0–T39.9, N14.0
4-Aminophenol derivatives	Paracetamol	T39.1
Antiepileptic, sedative-hypnotic and antiparkinsonism drugs	Barbiturates, Pregabalin, Benzodiazepines, Ketamine	F13.0*–F13.9*, T41.2*, T42.0–T42.8
Benzodiazepines	Benzodiazepines	T42.4
Cannabinoids	Cannabis, Cannabis derivatives	F12.0–F12.9, T40.7
Hallucinogens	LSD	F16.0*–F16.9*, T40.8–T40.9
Cocaine		F14.0–F14.9, T40.5
Amphetamine-type stimulants	Methamphetamine, MDMA, Caffeine	F15.0*–F15.9*, T43.6*
Antidepressants	Sertraline, Citalopram, Venlafaxine, Fluoxetine, Mirtazepine, Fluvoxamine, Paroxetine, Duloxetine	F55.0, T43.0–T43.2
Antipsychotics and neuroleptics	Quetiapine, Olanzapine, Risperidone	T43.3–T43.5
Volatile solvents	Petroleum products, nitrogen oxides	F18.0–F18.9, T52.0–T52.9, T53.0–T53.9, T59.0, T59.8
Multiple drug use		F19.0–F19.9

*ICD-10-AM code starting with the characters

Principal diagnosis and external cause

The following ICD-10-AM codes were used to examine trends in hospitalisations related to drug poisoning and mental and behavioural disorders due to substance use from 1999-00 to 2020-21.

Diagnosis category	ICD-10-AM
Poisoning	T40.0-T40.9, T39.0-T39.9, T42.0-T42.8, T43.0-T43.6, T52.0-T52.9, T53.0-T53.9, T59.0, T59.8
Mental and behavioural disorder due to substance use	F11-F16, F18 F19
Acute intoxication	Fxx.0*
Harmful use	Fxx.1*
Dependence	Fxx.2*
Withdrawal	Fxx.3*, Fxx.4*
Drug-induced psychotic disorder	Fxx.5*, Fxx.7*
Other	Fxx.6*, Fxx.8*, Fxx.9*

* Codes starting with 'F' and with the third digit as specified

External cause	ICD-10-AM
Unintentional	X40-X49
Intentional	X60-X69
Undetermined	Y10-Y19
Other	V00-Y98 excluding X40-X49, X60-X69, Y10-Y19, Y92, Y93

Opioid-related hospitalisations

The following ICD-10-AM codes were used to examine trends in opioid-related hospitalisations from 1999-00 to 2020-21.

Diagnosis category	ICD-10 diagnosis	ICD-10-AM
	<i>Poisoning by narcotics and psychodysleptics:</i>	
Poisoning	Opium poisoning	T40.0
Poisoning	Heroin poisoning	T40.1
Poisoning	Natural and semi-synthetic	T40.2
Poisoning	Methadone poisoning	T40.3
Poisoning	Other synthetic narcotics poisoning	T40.4
Poisoning	Other and unspecified narcotics poisoning	T40.6
Mental and behavioural disorder due to substance use	<i>Mental and behavioural disorders due to use of opioids:</i>	
Acute intoxication	Acute intoxication	F11.0
Harmful use	Harmful use	F11.1
Dependence	Dependence syndrome	F11.2
Withdrawal	Withdrawal state	F11.3
Withdrawal	Withdrawal state with delirium	F11.4
Drug-induced psychotic disorder	Psychotic disorder	F11.5
Other	Amnesic syndrome	F11.6
Drug-induced psychotic disorder	Residual and late-onset psychotic disorder	F11.7
Other	Other mental and behavioural disorders	F11.8
Other	Unspecified mental and behavioural disorder	F11.9

Non-opioid analgesic-related hospitalisations

The following ICD-10-AM codes were used to examine trends in non-opioid analgesic-related hospitalisations from 1999-00 to 2020-21.

Diagnosis category	ICD-10 diagnosis	ICD-10-AM
Harmful use	Harmful use of nondependence-producing substances: analgesics	F55.2
	<i>Poisoning by nonopioid analgesics, antipyretics and antirheumatics:</i>	
Poisoning	Salicylates	T39.0
Poisoning	4-Aminophenol derivatives	T39.1
Poisoning	Other nonsteroidal anti-inflammatory drugs [NSAID]	T39.3
Poisoning	Antirheumatics, not elsewhere classified	T39.4
Poisoning	Other nonopioid analgesics and antipyretics, not elsewhere classified	T39.8
Poisoning	Nonopioid analgesic, antipyretic and antirheumatic, unspecified	T39.9
Other	Drug- and heavy-metal-induced tubulo-interstitial and tubular conditions: analgesic nephropathy	N14.0

Antiepileptic, sedative-hypnotic and antiparkinsonism drug-related hospitalisations

The following ICD-10-AM codes were used to examine trends in antiepileptic, sedative-hypnotic and antiparkinsonism drug-related hospitalisations from 1999-00 to 2020-21.

Diagnosis category	ICD-10 diagnosis	ICD-10-AM
Poisoning	Poisoning by other and unspecified general anaesthetics	T41.2*
	<i>Poisoning by antiepileptic, sedative-hypnotic and antiparkinsonism drugs:</i>	
Poisoning	Hydantoin derivatives	T42.0
Poisoning	Iminostilbenes	T42.1
Poisoning	Succinimides and oxazolidinediones	T42.2
Poisoning	Barbiturates	T42.3
Poisoning	Benzodiazepines	T42.4
Poisoning	Mixed antiepileptics, not elsewhere classified	T42.5
Poisoning	Other antiepileptic and sedative-hypnotic drugs	T42.6
Poisoning	Antiepileptic and sedative-hypnotic drugs, unspecified	T42.7
Poisoning	Antiparkinsonism drugs and other central muscle-tone depressants	T42.8
Mental and behavioural disorder due to substance use	<i>Mental and behavioural disorders due to use of sedatives or hypnotics:</i>	F13.0*-F13.9*
Acute intoxication	Acute intoxication	F13.0*
Harmful use	Harmful use	F13.1*
Dependence	Dependence syndrome	F13.2*
Withdrawal	Withdrawal state	F13.3*
Withdrawal	Withdrawal state with delirium	F13.4*
Drug-induced psychotic disorder	Psychotic disorder	F13.5*
Other	Amnesic syndrome	F13.6*
Drug-induced psychotic disorder	Residual and late-onset psychotic disorder	F13.7*
Other	Other mental and behavioural disorders	F13.8*
Other	Unspecified mental and behavioural disorder	F13.9*

*ICD-10-AM code starting with the characters

Cannabinoid-related hospitalisations

The following ICD-10-AM codes were used to examine trends in cannabinoid-related hospitalisations from 1999-00 to 2020-21.

Diagnosis category	ICD-10 diagnosis	ICD-10-AM
Poisoning	Poisoning by narcotics and psychodysleptics: Cannabis (derivatives)	T40.7
Mental and behavioural disorder due to substance use	<i>Mental and behavioural disorders due to use of cannabinoids:</i>	F12.0-F12.9
Acute intoxication	Acute intoxication	F12.0
Harmful use	Harmful use	F12.1
Dependence	Dependence syndrome	F12.2
Withdrawal	Withdrawal state	F12.3
Withdrawal	Withdrawal state with delirium	F12.4
Drug-induced psychotic disorder	Psychotic disorder	F12.5
Other	Amnesic syndrome	F12.6
Drug-induced psychotic disorder	Residual and late-onset psychotic disorder	F12.7
Other	Other mental and behavioural disorders	F12.8
Other	Unspecified mental and behavioural disorder	F12.9

Hallucinogen-related hospital separations

The following ICD-10-AM codes were used to examine trends in hallucinogen-related hospitalisations from 1999-00 to 2020-21.

Diagnosis category	ICD-10 diagnosis	ICD-10-AM
	<i>Poisoning by narcotics and psychodysleptics [hallucinogens]:</i>	
Poisoning	Lysergide [LSD]	T40.8
Poisoning	Other and unspecified psychodysleptics [hallucinogens]	T40.9
Mental and behavioural disorder due to substance use	<i>Mental and behavioural disorders due to use of hallucinogens:</i>	F16.0*-F16.9*
Acute intoxication	Acute intoxication	F16.0*
Harmful use	Harmful use	F16.1*
Dependence	Dependence syndrome	F16.2*
Withdrawal	Withdrawal state	F16.3*
Withdrawal	Withdrawal state with delirium	F16.4*
Drug-induced psychotic disorder	Psychotic disorder	F16.5*
Other	Amnesic syndrome	F16.6*
Drug-induced psychotic disorder	Residual and late-onset psychotic disorder	F16.7*
Other	Other mental and behavioural disorders	F16.8*
Other	Unspecified mental and behavioural disorder	F16.9*

*ICD-10-AM code starting with the characters

Cocaine-related hospitalisations

The following ICD-10-AM codes were used to examine trends in cocaine-related hospitalisations from 1999-00 to 2020-21.

Diagnosis category	ICD-10 diagnosis	ICD-10-AM
Poisoning	Poisoning by narcotics and psychodysleptics: Cocaine	T40.5
Mental and behavioural disorder due to substance use	<i>Mental and behavioural disorders due to use of cocaine:</i>	F14.0-F14.9
Acute intoxication	Acute intoxication	F14.0
Harmful use	Harmful use	F14.1
Dependence	Dependence syndrome	F14.2
Withdrawal	Withdrawal state	F14.3
Withdrawal	Withdrawal state with delirium	F14.4
Drug-induced psychotic disorder	Psychotic disorder	F14.5
Other	Amnesic syndrome	F14.6
Drug-induced psychotic disorder	Residual and late-onset psychotic disorder	F14.7
Other	Other mental and behavioural disorders	F14.8
Other	Unspecified mental and behavioural disorder	F14.9

Amphetamine-type stimulant-related hospitalisations

The following ICD-10-AM codes were used to examine trends in amphetamine-type stimulant-related hospitalisations from 1999-00 to 2020-21.

Diagnosis category	ICD-10 diagnosis	ICD-10-AM
Poisoning	Poisoning by psychostimulants (excluding cocaine)	T43.6*
Mental and behavioural disorder due to substance use	<i>Mental and behavioural disorders due to use of other stimulants, including caffeine:</i>	F15.0*-F15.9*
Acute intoxication	Acute intoxication	F15.0*
Harmful use	Harmful use	F15.1*
Dependence	Dependence syndrome	F15.2*
Withdrawal	Withdrawal state	F15.3*
Withdrawal	Withdrawal state with delirium	F15.4*
Drug-induced psychotic disorder	Psychotic disorder	F15.5*
Other	Amnesic syndrome	F15.6*
Drug-induced psychotic disorder	Residual and late-onset psychotic disorder	F15.7*
Other	Other mental and behavioural disorders	F15.8*
Other	Unspecified mental and behavioural disorder	F15.9*

*ICD-10-AM code starting with the characters

Antidepressant, antipsychotic and neuroleptic-related hospitalisations

The following ICD-10-AM codes were used to examine trends in antidepressant, antipsychotic and neuroleptic-related hospitalisations from 1999-00 to 2020-21.

Diagnosis category	ICD-10 diagnosis	ICD-10-AM
Antidepressants		
Harmful use	Harmful use of nondependence-producing substances: antidepressants	F55.0
<i>Poisoning by psychotropic drugs, not elsewhere classified:</i>		
Poisoning	Tricyclic and tetracyclic antidepressants	T43.0
Poisoning	Monoamine-oxidase-inhibitor antidepressants	T43.1
Poisoning	Other and unspecified antidepressants	T43.2
Antipsychotics and neuroleptics		
<i>Poisoning by psychotropic drugs, not elsewhere classified:</i>		
Poisoning	Phenothiazine antipsychotics and neuroleptics	T43.3
Poisoning	Butyrophenone and thioxanthene neuroleptics	T43.4
Poisoning	Other and unspecified antipsychotics and neuroleptics	T43.5

Volatile solvent-related hospitalisations

The following ICD-10-AM codes were used to examine trends in volatile solvent-related hospitalisations from 1999-00 to 2020-21.

Diagnosis category	ICD-10 diagnosis	ICD-10-AM
	<i>Toxic effect of organic solvents:</i>	
Poisoning	Petroleum products	T52.0
Poisoning	Benzene	T52.1
Poisoning	Homologues of benzene	T52.2
Poisoning	Glycols	T52.3
Poisoning	Ketones	T52.4
Poisoning	Other organic solvents	T52.8
Poisoning	Organic solvent, unspecified	T52.9
	<i>Toxic effect of halogen derivatives of aliphatic and aromatic hydrocarbons:</i>	
Poisoning	Carbon tetrachloride	T53.0
Poisoning	Chloroform	T53.1
Poisoning	Trichloroethylene	T53.2
Poisoning	Tetrachloroethylene	T53.3
Poisoning	Dichloromethane	T53.4
Poisoning	Chlorofluorocarbons	T53.5
Poisoning	Other halogen derivatives of aliphatic hydrocarbons	T53.6
Poisoning	Other halogen derivatives of aromatic hydrocarbons	T53.7
Poisoning	Halogen derivative of aliphatic and aromatic hydrocarbons, unspecified	T53.9
	<i>Toxic effect of other gases, fumes and vapours:</i>	
Poisoning	Nitrogen oxides	T59.0
Poisoning	Other specified gases, fumes and vapours	T59.8
Mental and behavioural disorder due to substance use	<i>Mental and behavioural disorders due to use of volatile solvents:</i>	F18.0-F18.9
Acute intoxication	Acute intoxication	F18.0
Harmful use	Harmful use	F18.1
Dependence	Dependence syndrome	F18.2
Withdrawal	Withdrawal state	F18.3
Withdrawal	Withdrawal state with delirium	F18.4
Drug-induced psychotic disorder	Psychotic disorder	F18.5
Other	Amnesic syndrome	F18.6
Drug-induced psychotic disorder	Residual and late-onset psychotic disorder	F18.7
Other	Other mental and behavioural disorders	F18.8
Other	Unspecified mental and behavioural disorder	F18.9

Multiple drug use-related hospitalisations

The following ICD-10-AM codes were used to examine trends in multiple drug use-related hospitalisations from 1999-00 to 2020-21.

Diagnosis category	ICD-10 diagnosis	ICD-10-AM
Mental and behavioural disorder due to substance use	<i>Mental and behavioural disorders due to multiple drug use and use of psychoactive substances:</i>	F19.0-F19.9
Acute intoxication	Acute intoxication	F19.0
Harmful use	Harmful use	F19.1
Dependence	Dependence syndrome	F19.2
Withdrawal	Withdrawal state	F19.3
Withdrawal	Withdrawal state with delirium	F19.4
Drug-induced psychotic disorder	Psychotic disorder	F19.5
Other	Amnesic syndrome	F19.6
Drug-induced psychotic disorder	Residual and late-onset psychotic disorder	F19.7
Other	Other mental and behavioural disorders	F19.8
Other	Unspecified mental and behavioural disorder	F19.9