


There were 350 registered overdose and other drug-induced deaths (excluding alcohol and tobacco) in [Queensland](#) in 2021, which is equivalent to 1.0% of all registered deaths in this jurisdiction.


The population rate increased from 4.4 in 2002 to 7.4 in 2019 but peaked at 8.5 deaths per 100,000 people in 2015. The preliminary age-standardised rate of drug-induced deaths was 6.7 deaths per 100,000 people in 2021 ([Figure 1](#)). This rate was not statistically different from the estimated rate in 2020 (7.1 deaths per 100,000 people), noting that estimates for 2020 and 2021 are subject to revision and may increase (Table 1).

Sex

 In 2021, males accounted for 59% (205 deaths) of drug-induced deaths. The rate of drug-induced deaths was also higher among males than females (8.2 versus 5.2 deaths per 100,000 people, respectively). Analyses did not indicate a statistically significant difference between 2020 and 2021 in the estimated rates for males or females (Table 1).

Age

In 2021, drug-induced deaths were most common among the 35-44 age group (27%, 93 deaths). The rate was also highest in the 35-44 age group (13 deaths per 100,000 people), and in the 85 and over age group (13 deaths per 100,000 people).



Analyses did not indicate a statistically significant difference in the estimated rates between 2020 and 2021 for any of the age groups (Table 2).

Remoteness Area of Usual Residence

The greatest proportion of drug-induced deaths in 2021 occurred among people residing in major city areas (67%, 235 deaths), and the highest rate was also observed among people in major city areas (7.0 deaths per 100,000 people), followed by inner regional areas (6.5 deaths per 100,000 people).

There was no clear historical trend observed in the rate of drug-induced deaths for major city versus regional and remote areas of Queensland. The 2021 rates were comparable to the rates observed in 2020 (Table 3).

Intent of Drug Overdose Deaths

In 2021, 97% (339 deaths) of drug-induced deaths were due to [overdose](#). Unintentional drug overdose accounted for 60% (203 deaths) and intentional drug overdose for 35% (120 deaths) of these deaths in 2021. This profile was broadly consistent over time. Comparison of preliminary rates did not suggest a significant change between 2020 and 2021 (Table 4).

Place of Occurrence



In 2021, the location of the incident underlying death was coded as home for the majority (77%, 270 deaths) of drug-induced deaths.

Drug Involvement

In Queensland, the three most common drug types involved in drug overdose deaths in 2021 were:

- **opioids** (3.8 deaths per 100,000 people, 193 deaths),
- **antiepileptic, sedative-hypnotic and anti-parkinsonism drugs** (3.7 deaths per 100,000 people, 191 deaths),

- **antidepressants** (2.5 deaths per 100,000 people, 131 deaths) ([Figure 2](#)).

Comparison of estimated rates of drug overdose deaths in Queensland did not identify a significant change in rates of drug involvement between 2020 and 2021, noting that these are subject to revision and may increase (Table 5).

Figure 1. Age-standardised rate per 100,000 people of drug-induced deaths, by sex, Queensland, 2002-2021.





Figure 2. Age-standardised rate per 100,000 people of drug overdose deaths, by drug class, Queensland, 2002-2021.

Note: Deaths where conditions related to alcohol or tobacco comprised the underlying cause of death are not captured here.

Causes of death data for 2020 and 2021 are not final and thus are subject to further revision. The symbol 'o' indicates revised estimates and 'x' preliminary estimates.

Age-standardised rates were not calculated if the number of deaths was less than or equal to 10 (please refer to our [methods](#) document for details). Suppressed data are visible as gaps in the data series.

Table 1. Age-standardised rate per 100,000 people of drug-induced deaths in Queensland in 2020 and 2021, and average percent change (APC) for difference between 2021 and 2020 (with 95% confidence intervals), by sex

Sex	Rate in 2020 (95% CI)	Rate in 2021 (95% CI)	APC (95% CI)
Female	4.8 (4, 5.7)	5.2 (4.4, 6.2)	9.2 (-14, 39)
Male	9.5 (8.3, 11)	8.2 (7.1, 9.4)	-13 (-28, 4.9)
Total	7.1 (6.4, 7.9)	6.7 (6, 7.5)	-5.3 (-18, 9.9)

Note: Deaths where conditions related to alcohol or tobacco comprised the underlying cause of death are not captured here. Causes of death data for 2020 and 2021 are preliminary and thus are subject to further revision. 95% confidence intervals for the age-standardised rate and average percent change are shown in brackets. Please refer to our [methods](#) document on 'Presentation of results' for interpretation of average percent change. Please also refer to our [methods](#) document on 'Data source' and 'Coding of deaths' for details on the data used.

Table 2. Crude rate per 100,000 people of drug-induced deaths in Queensland in 2020 and 2021, and average percent change (APC) for difference between 2021 and 2020 (with 95% confidence intervals), by age

Age	Rate in 2020 (95% CI)	Rate in 2021 (95% CI)	APC (95% CI)
15-24	3.5 (2.2, 5.3)	3.5 (2.2, 5.3)	0.91 (-46, 88)
25-34	8.8 (6.8, 11)	6.5 (4.8, 8.7)	-26 (-50, 9.9)
35-44	15 (12, 18)	13 (11, 16)	-8.7 (-32, 22)

45-54	11 (8.4, 13)	11 (9.1, 14)	7.3 (-23, 50)
55-64	9.7 (7.4, 12)	9.4 (7.1, 12)	-2.8 (-34, 42)
65-74	5 (3.2, 7.4)	4.2 (2.6, 6.4)	-15 (-55, 59)
75-84	6.3 (3.6, 10)	6.3 (3.7, 10)	0.26 (-52, 112)
85+	–	13 (6.5, 22)	–

Note: Deaths where conditions related to alcohol or tobacco comprised the underlying cause of death are not captured here. Causes of death data for 2020 and 2021 are preliminary and thus are subject to further revision. 95% confidence intervals for the crude rate and average percent change are shown in brackets. Please refer to our [methods](#) document on 'Presentation of results' for interpretation of average percent change. The estimates for the 0-14 years age group are not presented due to sensitivity of the data. Please also refer to our [methods](#) document on 'Data source' and 'Coding of deaths' for details on the data used.

Table 3. Age-standardised rate per 100,000 people of drug-induced deaths in Queensland in 2020 and 2021, and average percent change (APC) for difference between 2021 and 2020 (with 95% confidence intervals), by remoteness area

Remoteness	Rate in 2020 (95% CI)	Rate in 2021 (95% CI)	APC (95% CI)
Major Cities	7.3 (6.4, 8.3)	7 (6.1, 7.9)	-4.6 (-20, 14)
Regional and Remote	6.5 (5.4, 7.9)	5.9 (4.8, 7.1)	-9.8 (-31, 18)

Note: Deaths where conditions related to alcohol or tobacco comprised the underlying cause of death are not captured here. Causes of death data for 2020 and 2021 are preliminary and thus are subject to further revision. 95% confidence intervals for the age-standardised rate and average percent change are shown in brackets. Please refer to our [methods](#) document on 'Presentation of results' for interpretation of average percent change. Please also refer to our [methods](#) document on 'Data source' and 'Coding of deaths' for details on the data used.

Table 4. Age-standardised rate per 100,000 people of overdose deaths in Queensland in 2020 and 2021, and average percent change (APC) for difference between 2021 and 2020 (with 95% confidence intervals), by intent

Intent	Rate in 2020 (95% CI)	Rate in 2021 (95% CI)	APC (95% CI)
Unintentional	4.5 (4, 5.2)	4.1 (3.5, 4.7)	-9.7 (-25, 9.3)
Intentional	1.9 (1.5, 2.3)	2.1 (1.8, 2.6)	13 (-14, 47)

Note: Deaths where conditions related to alcohol or tobacco comprised the underlying cause of death are not captured here. Causes of death data for 2020 and 2021 are preliminary and thus are subject to further revision. 95% confidence intervals for the age-standardised rate and average percent change are shown in brackets. Please refer to our [methods](#) document on 'Presentation of results' for interpretation of average percent change. Please also refer to our [methods](#) document on 'Data source' and 'Coding of deaths' for details on the data used.

Table 5. Age-standardised rate per 100,000 people of overdose deaths in Queensland in 2020 and 2021, and average percent change (APC) for difference between 2021 and 2020 (with 95% confidence intervals), by drugs involved

Drug	Rate in 2020 (95% CI)	Rate in 2021 (95% CI)	APC (95% CI)
Opioids	4.5 (3.9, 5.1)	3.8 (3.3, 4.4)	-16 (-31, 2.2)
Antiepileptic, sedative-hypnotic & antiparkinsonism drugs	4.6 (4.0, 5.2)	3.7 (3.2, 4.2)	-20 (-34, -2.4)
Antidepressants	2.9 (2.5, 3.4)	2.5 (2, 2.9)	-16 (-34, 6.9)
Amphetamine-type stimulants	2.3 (1.9, 2.8)	1.4 (1.1, 1.8)	-40 (-56, -18)
Antipsychotics & neuroleptics	1.5 (1.2, 1.9)	1.4 (1.1, 1.7)	-11 (-36, 23)
Alcohol	1.3 (1.0, 1.7)	1.2 (0.89, 1.5)	-13 (-38, 24)
Non-opioid analgesics	0.93 (0.69, 1.2)	0.81 (0.59, 1.1)	-13 (-43, 32)
Cannabinoids	0.6 (0.4, 0.87)	0.27 (0.14, 0.46)	-56 (-77, -14)
Cocaine	0.29 (0.16, 0.49)	0.25 (0.13, 0.43)	-16 (-61, 82)

Note: Deaths where conditions related to alcohol or tobacco comprised the underlying cause of death are not captured here. Causes of death data for 2020 and 2021 are preliminary and thus are subject to further revision. 95% confidence intervals for the age-standardised rate and average percent

change (APC) are shown in brackets. Please refer to our [methods](#) document on 'Presentation of results' for interpretation of average percent change. Please also refer to our [methods](#) document on 'Data source' and 'Coding of deaths' for details on the data used.

ISSN XXXX ©NDARC, UNSW SYDNEY 2023

This report was prepared by researchers from the National Drug and Alcohol Research Centre for the Drug Trends program. The Drug Trends program is coordinated by the National Drug and Alcohol Research Centre, UNSW Sydney and undertaken in partnership with the Burnet Institute, National Drug Research Institute, University of Queensland, and University of Tasmania.

This work is copyright. You may download, display, print and reproduce this material in unaltered form only (retaining this notice) for your personal, non-commercial use or use within your organisation. All other rights are reserved. Requests and enquiries concerning reproduction and rights should be addressed to NDARC, UNSW Sydney, NSW 2052, Australia.

Recommended citation: Chrzanowska A, Man N, Akhurst J, Sutherland R, Degenhardt L, Peacock A. Trends in overdose and other drug-induced deaths in Australia, 2002-2021. Sydney: National Drug and Alcohol Research Centre, UNSW Sydney; 2023. Available from: <http://doi.org/10.26190/m2gs-z325>

Please note that as with all statistical reports, there is the potential for minor revisions to data in this report. Please refer to the online version at [Drug Trends](#).

Please contact the Drug Trends team with any queries regarding this publication: drugtrends@unsw.edu.au.

Funding

The Drug Trends program is funded by the Australian Government Department of Health and Aged Care under the Drug and Alcohol Program.

Data source

We acknowledge all state and territory Registries of Births, Deaths and Marriages, the Coroners and the National Coronial Information System (NCIS) for enabling Cause of Death Unit Record File (COD URF) data to be used for this publication.

Acknowledgements

We wish to acknowledge Lauren Moran and the team at the Australian Bureau of Statistics for their assistance with the data and ICD-10 coding practices to ensure rigorous, comprehensive, and consistent reporting on drug-induced deaths in Australia.

We would like to acknowledge the contribution of those who have been involved in past reporting on drug-induced deaths by Drug Trends, specifically: A/Prof Timothy Dobbins, Dr Amanda Roxburgh, and A/Prof Lucinda Burns.

We acknowledge the traditional custodians of the land on which the work for this report was undertaken. We pay respect to Elders past, present, and emerging.

Related Links

- For interactive data visualisations accompanying this report, go to: https://drugtrends.shinyapps.io/Deaths_2021
- For full details of the methods underpinning this report, go to: <https://ndarc.med.unsw.edu.au/resource-analytics/trends-drug-induced-deaths-australia-2002-2021>
- For other Drug Trends publications on drug-related hospitalisations and drug-induced deaths in Australia, go to: <https://ndarc.med.unsw.edu.au/project/national-illicit-drug-indicators-project-nidip>
- For more information on NDARC research, go to: <http://ndarc.med.unsw.edu.au/>

- For more information about the ABS, go to: <http://www.abs.gov.au>
- For more information on ICD coding go to: <http://www.who.int/classifications/icd/en/>
- For more information on the Remoteness Areas Structure within the Australian Statistical Geography Standard (ASGS), go to: <https://www.abs.gov.au/ausstats/abs@.nsf/mf/1270.0.55.005>
- For more research from the Drug Trends program and to subscribe to our newsletter, go to: <https://ndarc.med.unsw.edu.au/program/drug-trends>
- For details on the collection, organisation and interpretation of NCIS data, go to: <https://www.ncis.org.au/about-the-data/explanatory-notes/>
- For statistics about case closure statistics in NCIS, go to: <https://www.ncis.org.au/about-the-data/operational-statistics/>